

Vogel, J. T., Somers, M. J. and Venter, J. A. 2018. The foraging ecology of reintroduced African wild dog in small protected areas. – Wildlife Biology 2018: wlb.00424

Appendix 1

Table A1. The corrected prey biomass (kg) for pooled wild dog kills.

Species	Kills		Uncorrected mass (kg)			Corrected mass (kg)	
	n	Male	Female	Subadult	Juvenile	Kg	%
Buffalo	3	650.00	520.00	364.00	156.00	313.56	1.41
Bushbuck	8	60.00	36.00	25.20	10.80	334.80	1.50
Bushpig	1	60.00		42.00	18.00	45.00	0.20
Cane Rat	1	3.81		2.67	1.14	2.83	0.01
Common Reedbuck	11	70.00	51.00	35.70	15.30	527.63	2.37
Grey Duiker	14	17.00	21.00	14.70	6.30	228.15	1.03
Impala	213	60.00	45.00	31.50	13.50	6742.80	30.30
Kudu	26	225.00	155.00	108.50	46.50	2283.40	10.26
Mountain Reedbuck	3	30.00	24.00	16.80	7.20	75.60	0.34
Nyala	190	108.00	62.00	43.40	18.60	8245.81	37.05
Red Duiker	18	12.00	12.00	8.40	3.60	187.16	0.84
Steenbuck	2	11.00	11.30	7.91	3.39	20.07	0.09
Warthog	37	80.00	60.00	42.00	18.00	1279.20	5.75
Waterbuck	4	260.00	180.00	126.00	54.00	429.10	1.93
Wildebeest	20	250.00	180.00	126.00	54.00	1410.67	6.34
Zebra	2	320.00		224.00	96.00	128.64	0.58
	553					22254.41	100.00